

Doubly charged Higgs in a 3-3-1 model at the CERN LHC
Prospects for Charged Higgs Discovery at Colliders

16-19 September 2008

Uppsala, Sweden

J. E. Cieza Montalvo¹, Nelson V. Cortez*, M. D. Tonasse²

¹*Rio de Janeiro State University,*

²*Sao Paulo State University, Brazil*

**, Speaker*

Abstract

Doubly charged Higgs bosons are very important in particle physics nowadays because they can give mass to neutrinos through seesaw mechanisms. In this work we present the results in the searching for doubly charged Higgs at the CERN LHC for a 3-3-1 model.

CERN LHC must bring great news in physics fields in the next years. In special, the discovery of Higgs boson and extra-dimensions occupy a very important part of the physicists' mind worldwide. Higgs bosons, for instance, can signalize to new models where these particles can be singly and/or doubly charged. There is a big interest in doubly charged Higgs bosons (DCHBs) due to seesaw mechanisms which can give mass to neutrinos, as for instance in Ref. [1].

Very interesting results were obtained recently in the searching for DCHBs with lepton flavor violating decays involving tau leptons using Left-Right Symmetric model [2]. However, an important class of models was raised through an efficient chiral expansion of the Standard Model(SM) in the electroweak sector, i.e. the full gauge symmetry passed from $SU(3)_C \otimes SU(2)_W \otimes U(1)_Y$ to $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$, where $SU(2)_W \subset SU(3)_W$. For short it has been named 3-3-1 Model, see Ref. [3]. A very recent exciting manuscript has suggested that the observed excess of muons in studies in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV arises from the decay of doubly-charged bosons in particles predicted in the 3-3-1 model [4].

The authors of this present work have worked with different kind of particle colliders searching for DCHBs as can be seen in Refs. [5]. They have worked with a 3-3-1 model which has the simplest scalar sector in this class of models [6]. An overview of it is presented below with results for the CERN LHC.

THE 3-3-1 MODEL: The underlying electroweak symmetry group is $SU(3)_L \otimes U(1)_N$, where N is the quantum number of the $U(1)$ group. Therefore, the left-handed lepton matter content is $\left(\nu'_a \ \ell'_a \ L'_a \right)_L^T$ transforming as $(\mathbf{3}, 0)$, where $a = e, \mu, \tau$ is a family index (we are using primes for the interaction eigenstates). L'_{aL} are lepton fields which can be the charge conjugates ℓ'^{+}_{aR} [3] or the antineutrinos ν'^{+}_{La} or heavy leptons P'^{+}_{aL} ($P'^{+}_{aL} = E'^{+}_L, M'^{+}_L, T'^{+}_L$) [6]. This model has the charge operator given by $\frac{Q}{e} = \frac{1}{2} (\lambda_3 - \sqrt{3}\lambda_8) + N$, where λ_3 and λ_8 are the diagonal Gell-Mann matrices and e is the elementary electric charge. The right-handed charged leptons are introduced in singlet representation of $SU(3)_L$ as $\ell'^{-}_{aR} \sim (\mathbf{1}, -1)$ and $P'^{+}_{aR} \sim (\mathbf{1}, 1)$.

The model has three scalar triplets in the minimal scalar sector:

$$\eta = \begin{pmatrix} \eta^0 \\ \eta_1^- \\ \eta_2^+ \end{pmatrix} \sim (\mathbf{3}, 0), \quad \rho = \begin{pmatrix} \rho^+ \\ \rho^0 \\ \rho^{++} \end{pmatrix} \sim (\mathbf{3}, 1), \quad \chi = \begin{pmatrix} \chi^- \\ \chi^{--} \\ \chi^0 \end{pmatrix} \sim (\mathbf{3}, -1). \quad (1a)$$

The most general, gauge invariant and renormalizable Higgs potential, which conserves the

leptobaryon number [7], is

$$\begin{aligned}
V(\eta, \rho, \chi) = & \mu_1^2 \eta^\dagger \eta + \mu_2^2 \rho^\dagger \rho + \mu_3^2 \chi^\dagger \chi + \lambda_1 (\eta^\dagger \eta)^2 + \lambda_2 (\rho^\dagger \rho)^2 + \lambda_3 (\chi^\dagger \chi)^2 + \\
& + (\eta^\dagger \eta) [\lambda_4 (\rho^\dagger \rho) + \lambda_5 (\chi^\dagger \chi)] + \lambda_6 (\rho^\dagger \rho) (\chi^\dagger \chi) + \lambda_7 (\rho^\dagger \eta) (\eta^\dagger \rho) + \\
& + \lambda_8 (\chi^\dagger \eta) (\eta^\dagger \chi) + \lambda_9 (\rho^\dagger \chi) (\chi^\dagger \rho) + \frac{1}{2} (f \epsilon^{ijk} \eta_i \rho_j \chi_k + \text{H. c.})
\end{aligned} \tag{2}$$

The neutral components of the scalars triplets (1a) develop non zero vacuum expectation values (VEV's) $\langle \eta^0 \rangle = v_\eta$, $\langle \rho^0 \rangle = v_\rho$ and $\langle \chi^0 \rangle = v_\chi$, with $v_\eta^2 + v_\rho^2 = v_W^2 = (246 \text{ GeV})^2$. The pattern of symmetry breaking is $\text{SU}(3)_L \otimes \text{U}(1)_N \xrightarrow{\langle \chi \rangle} \text{SU}(2)_L \otimes \text{U}(1)_Y \xrightarrow{\langle \eta, \rho \rangle} \text{U}(1)_{\text{em}}$. Therefore, we can expect $v_\chi \gg v_\eta, v_\rho$. In the potential (2), f and μ_j ($j = 1, 2, 3$) are constants with dimension of mass and the λ_i ($i = 1, \dots, 9$) are adimensional constants. The procedures to obtain the masses of the Higgs bosons are showed in Ref. [8] under the conditions $v_\chi \approx -f$, leading a following results for the masses of the neutral physical scalars

$$m_{H_1^0}^2 \approx 4 \frac{\lambda_2 v_\rho^4 - 2\lambda_1 v_\eta^4}{v_\eta^2 - v_\rho^2}, \quad m_{H_2^0}^2 \approx \frac{v_W^2 v_\chi^2}{2v_\eta v_\rho}, \quad m_{H_3^0}^2 \approx -\lambda_3 v_\chi^2, \quad m_h^2 = -\frac{f v_\chi}{v_\eta v_\rho} \left[v_W^2 + \left(\frac{v_\eta v_\rho}{v_\chi} \right)^2 \right], \tag{3a}$$

for the singly charged ones

$$m_{\pm 1}^2 = \frac{v_W^2}{2v_\eta v_\rho} (f v_\chi - 2\lambda_7 v_\eta v_\rho), \quad m_{\pm 2}^2 = \frac{v_\eta^2 + v_\chi^2}{2v_\eta v_\chi} (f v_\rho - 2\lambda_8 v_\eta v_\chi), \tag{3b}$$

and for the doubly charged Higgs bosons

$$m_{\pm\pm}^2 = \frac{v_\rho^2 + v_\chi^2}{2v_\rho v_\chi} (f v_\eta - 2\lambda_9 v_\rho v_\chi). \tag{3c}$$

The total leptonic number is conserved [3] although that non standard field interactions violate leptonic number.

RESULTS

When we consider particular values for the adimensional parameter λ_9 in the Higgs potential (2) and for the vacuum expectation value v_χ [8] and with all constraints in the model used in this work, so the production and signals for the DCHBs in the 3-3-1 models at LHC collider are very significant. We have fixed the general bounds of the adimensional constants of the Higgs potential (2) as $-3 \leq \lambda_i \leq 3$ ($i = 1, \dots, 9$) to guarantee approximately the perturbative regime. When we consider $-f \approx v_\chi \gg v_\eta, v_\rho$, we have

$$\lambda_4 \approx 2 \frac{\lambda_2 v_\rho^2 - \lambda_1 v_\eta^2}{v_\eta^2 - v_\rho^2}, \quad \lambda_5 v_\eta^2 + 2\lambda_6 v_\rho^2 \approx -\frac{v_\eta v_\rho}{2} \tag{4}$$




FIG. 1: Parameter f as a function of mass of doubly charged Higgs boson for the parameter $\lambda_9 = -0.8$.




FIG. 2: Branching ratios for the doubly charged Higgs decays as a functions of $m_{H^{\pm\pm}}$ for $\lambda_9 = -0.8$ for the fermionic sector.




FIG. 3: Branching ratios for the doubly charged Higgs decays as a functions of $m_{H^{\pm\pm}}$ for $\lambda_9 = -0.8$ for the bosonic sector.




FIG. 4: Total cross section for the process $pp \rightarrow H^{--} H^{++}$ as a function of $m_{H^{\pm\pm}}$ for $\lambda_9 = -0.8$

as seen in Ref. [8].

A analysis of Eq. (3c) also shows that the lowest values for λ_9 and f , such to produce values for $m_{H^{\pm\pm}}$ between 100 GeV and 200 GeV, for $v_\chi = 1500$ GeV and $v_\rho = 195$ GeV are $\lambda_9 = -4.50 \times 10^{-3}$, $f = -0.23$ GeV and $\lambda_9 = -1.80 \times 10^{-2}$, $f = -0.92$ GeV. The values for the parameters λ_5 and λ_6 lead to the constraint $v_\eta > 40.5$ GeV, which can be seen as

TABLE I: Total number of events per year for DCHBs and with respect the branching ratios from $H^{\pm\pm} \rightarrow e^+P^+ + e^-P^-$ in function of m_H^{++} , $m_{H_2}^{\pm}$, m_h and v_χ , which their units are in GeV.

v_χ	m_H^{++}	$m_{H_2}^{\pm}$	m_h	Total events per year	Events per year for BR($H^{\pm\pm} \rightarrow e^+P^+ + e^-P^-$)
0.8	500	1163.7	2222.9	39×10^6	1.3×10^5
	700	1223.6	2052.2	2.2×10^6	3.9×10^3

following from the Eq. (4). In the approximation $-f \approx v_\chi$ it is appropriate to choose the parameter $-1.2 \leq \lambda_9 \leq -0.8$ for the mass of $m_{H^{\pm\pm}} = 500$ GeV. For simplicity we present here only the results for $\lambda_9 = -0.8$. The VEV's values v_η and v_ρ influence very little on the mass of $m_{H^{\pm\pm}}$, this happen because in the Eq. (3c) the v_η and v_ρ are related by the relation $v_\eta^2 + v_\rho^2 = (246 \text{ GeV})^2$ and v_ρ is smaller compared with v_χ .

We have chosen for the parameters, masses and the VEV, the following representative values: $\lambda_1 = -1.2$, $\lambda_2 = \lambda_3 = -\lambda_6 = \lambda_8 = -1$, $\lambda_4 = 2.98$, $\lambda_5 = -1.57$, $\lambda_7 = -2$ and $\lambda_9 = -0.8$, $v_\eta = 195$ GeV and $v_\chi = 1500$ GeV. Others particles, such as $J_{1,2,3}$, T , $U^{\pm\pm}$, Z' , $H_2^\pm$, $H_{2,3}^0$ and h^0 , do not take part in this process because of their masses are larger than the mass of the DCHBs. Observing the Fig. 1 we see that for the $v_\chi = 1000$ GeV we have the acceptable masses up to $m_{H^{\pm\pm}} \simeq 903$ GeV, for $v_\chi = 1500$ GeV we will have up to $m_{H^{\pm\pm}} \simeq 1346$ GeV and for $v_\chi = 1700$ GeV we have consequently up to $m_{H^{\pm\pm}} \simeq 1525$ GeV. Figs. 2 and 3, show the branching ratios for the Higgs decays, $H^{\pm\pm} \rightarrow \text{all}$. We have chosen $v_\chi = 1700$ GeV as the maximum value because it is constraint by the mass of the $m_{Z'} = (0.5 - 3)$ TeV, which is proportional to v_χ [3]

Considering that the expected integrated luminosity for the LHC will be of order of $3 \times 10^5 \text{ pb}^{-1}/\text{yr}$ then the statistics give a total of $\simeq 3.9 \times 10^6$ events per year for Drell-Yan process, if we take the mass of the Higgs boson $m_{H^{\pm\pm}} = 500$ GeV, $v_\chi = 1500$ GeV, $m_{H_2}^\pm = 1163.7$ GeV, $m_h = 2229.9$ GeV and $\lambda_9 = -0.8$. Considering that the signal for $H^{\pm\pm}$ production will be e^-P^- and e^+P^+ and taking into account that the branching ratios for both particles we would have approximately 1.3×10^5 events per year for Drell-Yan process. Taking now the mass of the Higgs boson $m_{H^{\pm\pm}} = 700$ GeV, $v_\chi = 1500$ GeV, $m_{H_2}^\pm = 1223.6$ GeV, $m_h = 2052.2$ GeV and the same value of λ_9 as above. We then have a total of $\simeq 2.2 \times 10^6$ events per year for Drell-Yan process. Fig. 4 shows the cross section for the process $pp \rightarrow H^{++}H^{--}$.

The main background for this signal, $p p \rightarrow H^{--}H^{++} \rightarrow e^-P^- (e^+P^+)$, could come from the process $p p \rightarrow Z Z$ and another small background from the $p p \rightarrow W^-W^+Z$, all these backgrounds can be eliminated (see Ref. [9]). Our studies have indicated the possibility of obtaining a clear signal of the doubly charged Higgs bosons, with a satisfactory number of events, in the next particle colliders and in a very special way to the CERN LHC, as can be seen in Table I.

-
- [1] N. V. Cortez and M. D. Tonasse, Phys. Rev. D **72**, 073005 (2005);
 - [2] T. Aaltonen *et al.* [The CDF Collaboration], Phys. Rev. Lett. **101**, 121801 (2008)
 - [3] F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992); P. H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992);
 - [4] P. H. Frampton, arXiv:0811.3943 [hep-ph].
 - [5] J. E. Cieza Montalvo, Nelson V. Cortez, J. Sá Borges and M. D. Tonasse, Nucl. Phys. B **756**, 1 (2006); J. E. Cieza Montalvo, N. V. Cortez and M. D. Tonasse, Phys. Rev. D **76**, 117703 (2007) ; J. E. Cieza Montalvo, Nelson V. Cortez, and M. D. Tonasse, Phys. Rev. D **77**, 095015 (2008); J. E. Cieza Montalvo, Nelson V. Cortez, and M. D. Tonasse, Phys. Rev. D **78**, 116003 (2008).
 - [6] V. Pleitez and M. D. Tonasse, Phys. Rev. D **48**, 2353 (1993); M. D. Tonasse, Phys. Lett. B **381**, 191 (1996)
 - [7] V. Pleitez and M. D. Tonasse, Phys. Rev. D **48**, 5274 (1993);
 - [8] M. D. Tonasse, Phys. Lett. B **381**, 191 (1996).
 - [9] J. E. Cieza Montalvo and P. P. Queiroz Filho, Phys. Rev. D **66**, 055003 (2002).